



**Composition of functions:** It will help if you think of functions as machines that take an input and produce an output. Consider two functions, f(x) and g(x), pictured below.



Let's suppose we apply the function g to a number, **and then** put that output into the function f. That is the idea of composition and is pictured below.



We will start our examples with a situation that helps justify why you would ever want to do this composition stuff.


expl 1: A popular umbrella manufacturer has this information.

We can calculate  $P(x) = 2x^2 - 3x$ , where P(x) represents the monthly profit of the company (in dollars) and x represents the number of umbrellas sold that month,  $x \ge 0$ .


We can calculate U(t) = 3t + 4, where U(t) represents the number of umbrellas sold per month and t represents the number of days it rains that month,  $0 \le t \le 31$ .

We are interested in how profit is related to the number of days of rain. Let's find the equation. But first, we'll investigate the situation with a specific example.

Specific example: Say we have had 12 days of rain this month. Find the number of umbrellas we can expect to sell and then the profit this should yield.



Notice how this was a two-step process. Composition of functions can shortcut these steps. I have illustrated this below.





Let's find the relationship between days of rain and profit.

We know  $P(x) = 2x^2 - 3x$  where x represents the number of umbrellas.

But U(t) also represents the number of umbrellas. So we can substitute U(t) in for x to find P(U(t)).

Do it now to make a new function that tells us the profit given the number of days of rain.






Check your answer by using your new function to calculate the profit when it rains 12 days this month. Does it match what you found earlier?

**Composition Notation:** We used the nested notation for the previous example. The following notations are equivalent.


$$(f \circ g)(x) = f \circ g = f(g(x))$$
  
Use the outputs of  $g$   
as the inputs of  $f$ .  
Pronounced  
"f of g of x"



b.) Find  $(f \circ f)(1)$ .


c.) Find  $(f \circ g)(x)$ . d.) Find  $(g \circ f)(x)$ .

**Domain of Composed Functions:** The domain of f(g(x)) is all real numbers x that are in the domain of g and such that g(x) is in the domain of f.



expl 4: Bob throws a rock into a still lake. The rock causes a circular ripple that gets bigger and bigger. The radius of the ripple is increasing at the rate of 2.6 feet per second. Answer the questions that follow.


a.) Find the function for the radius of the circular ripple at time t.



c.) Find the composed function  $(A \circ r)(t)$ . What, in words, does this tell us about the ripple?

## Worksheet: The Composition of Functions:

This worksheet provides practice in performing compositions. There is an application that will help explain why composition is so useful.



expl 6: Given the function h(x) below, find two functions f(x) and g(x) such that h(x) = f(g(x)).  $h(x) = \sqrt{4x^2 - 3}$ There are many answers.

expl 7: Let f(x) = 3x + 2 and  $g(x) = \frac{x-2}{3}$ . Find f(g(x)) and g(f(x)).

